

(19)



Eur päisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 0 712 935 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
22.05.1996 Bulletin 1996/21

(21) Application number: 95203153.2

(22) Date of filing: 17.11.1995

(51) Int. Cl.<sup>6</sup>: C12N 15/74, C12N 1/21,  
A23C 19/032, A23C 23/00,  
G01N 33/04

(84) Designated Contracting States:  
BE CH DE DK ES FR GB IE IT LI NL SE

(30) Priority: 18.11.1994 NL 9401934  
18.11.1994 NL 9401935

(71) Applicant: STICHTING NEDERLANDS INSTITUUT  
VOOR ZUIVELONDERZOEK  
NL-6710 BA Ede (NL)

(72) Inventors:  
• Kuipers, Oscar Paul  
NL-6713 MS Ede (NL)  
• De Vos, Willem Meindert  
NL-6721 SJ Bennekom (NL)

(74) Representative: de Bruijn, Leendert C. et al  
Nederlandsch Octrooibureau  
P.O. Box 29720  
NL-2502 LS Den Haag (NL)

**(54) Method for controlling the gene expression in lactic acid bacteria**

(57) The present invention relates to a method for the controlled expression, in a lactic acid bacterium, of a DNA fragment containing one or more genes coding for a desired characteristic, characterised in that

- the DNA fragment is under the control of a promoter for a microbial gene which codes for an antimicrobial peptide, and
- the gene/the genes are brought to expression on the DNA fragment by the addition of a suitable inducing factor for the transcription activation.


The promoter is preferably the *nisA* promoter from *Lactococcus lactis*.

The inducer is preferably acceptable for food products, and more preferentially is nisin or derivatives thereof.

The invention also relates to a method for the production of proteins or RNA, as well as to a method for the preparation of dairy products using the expression system according to the invention.

The invention finally relates to lactic acid bacteria and expression vectors for use in the method according to the invention.

fig -11



EP 0 712 935 A2

## Description

The present invention relates to a method for the controlled expression, in a lactic acid bacterium, of a DNA fragment which contains one or more genes which code for a desired characteristic.

Methods for the controlled expression of DNA fragments in lactic acid bacteria have already been described. However, these known systems are not satisfactory for various reasons.

For instance, J.M. Wells et al., Appl. Environm. Microbiol. 59 (1993) 3954-3959 describe a system for regulation of gene expression by an extracellular factor, in which system the heterologous T7 RNA polymerase gene is brought to controlled expression under the control of the *lac* promoter from the lactose operon of *L. lactis*. With this procedure the expression is induced by replacing glucose by lactose in the culture medium. However, this system - and comparable systems from the prior art in which the gene expression is induced by changing the sugar source in the culture medium - has a number of disadvantages, in particular in the case of large-scale preparation of, for example, foodstuffs, such as:

- The achievable degree of overexpression is restricted, usually to a factor of at most a few tens.
- The regulation is not strict/absolute, that is to say without the presence of the inducer some transcription of the regulated DNA fragment still takes place.
- The systems require the use of specific sugar sources, which can be expensive, such as, for example, xylose, which, moreover, have to be added in preponderant amounts.
- The sugar inducer can be added effectively only after removal of the sugar used beforehand.
- The presence of specific sugars in the end product is not always desirable, whilst the removal thereof - if this is even possible - is associated with additional production steps and additional costs.
- In some cases heterologous DNA must be introduced; for instance, the T7 RNA polymerase gene and promoter are used in the abovementioned system.

Lakshmidevi et al., Applied Environmental Microbiology, April 1990, p. 934-942 confirm the problems which prior to the present application existed in this specialist field:

"This program of starter improvement would be enhanced by the development of procedures enabling controlled expression of specific genes. To achieve this a detailed knowledge of the regulation of gene expression in these bacteria [*Lactococci*] is required. [...] However to our knowledge there are no published reports of detailed study on the regulation of gene expression in *Lactococci*."

Lakshmidevi himself describes the promoter of the bacteriophage Bk5-T which is effective in *Lactococcus*. However, this is not a bacterial promoter, nor a promoter which controls the expression of an antimicrobial peptide. Moreover, the bacteriophage Bk5-T DNA is heterologous with respect to lactic acid bacteria and it is also not possible to obtain a strict/absolute regulation of the expression.

In H. Israelsen et al.: "Environmentally regulated promoters in lactococci", Abstract 16, 4th Conference on Streptococcal genetics, Santa Fé, May 15-18 (1994) various induction systems are described for *L. lactis*, the gene expression being controlled by changing the temperature, growth phase, oxygen concentration and/or substrate components.

A temperature-sensitive regulation of the expression of the *dnaJ* gene of *L. lactis* has also been described (J. Bacteriol. 175 [1993] 1637-1644).

International Patent Application 94/00581 (Viagen Oy) describes a *Lactobacillus* expression system based on the gene sequences of the surface protein (sp) gene.

However, this is not a gene that codes for an antimicrobial peptide. Furthermore, the relevant promoter or the relevant operon is not induced by its own gene product.

Moreover, it is stated on page 6, lines 20-27, that "the *Lactobacillus* coat protein is apparently synthesized under a wide variety of culture conditions [...]. The use of this promoter provides an advantage when it is desired to increase the expressions of genes or operons of, for example, biosynthetic or degradative pathways which are otherwise tightly regulated".

The gene expression system described in said application therefore does not give a very stringent regulation, whilst the aim of the present invention is to obtain a very strict/absolute regulation.

D. Kirsch et al., Journal of Antibiotics, Part 2, No. 91, 1991, describe an assay for the detection of antibacterial antibiotics, wherein said antibiotics are added to microorganisms which contain fusions between promoters which can be initiated by antibiotics and structural genes for *E. coli*  $\beta$ -galactosidase.

The promoters mentioned in this context are the tetA/tetR promoter of transposon Tn10 induced by tetracycline, the promoter of *S. aureus* *erm* C. erythromycin-resistance gene induced by erythromycin and the promoter of *S. aureus* cat 86 chloramphenicol resistance gene induced by chloramphenicol.

However, these are promoters of resistance genes; such promoters can, of course, be induced by the addition of the relevant antibiotic. However, these antibiotics are not antimicrobial peptides, nor are they the gene product of the relevant resistance genes, nor are they food-grade.

International Patent Application 87/00203 (Columbia University) describes plasmids which contain promoters for bacteriocins such as colicin. These promoters can be used to regulate the expression in *E. coli*. However, regulation of the expression in lactic acid bacteria using this promoter is neither described nor suggested.

Also, these promoters cannot be induced by their own gene product, that is to say colicin itself and/or derivatives thereof. According to said application, the colicin synthesis can also be induced by a large number of different factors, such as factors in the culture medium, changes in the temperature, and, for example, mitomycin C. There is therefore no question of a strict/absolute regulation.

International Patent Application 92/04451 (Genesit Oy) describes, in general terms, vectors for promoters which are effective in *E. coli*, *B. subtilis*, *Lactococci* and *Lactobacillus*, chosen from the plasmids pKHT1734 and pKHT1736 described therein.

Neither the *nisA* promoter, nor any other *Lactococcus* promoter, the induction thereof or the use thereof for *Lactococci* is either described or suggested in said application.

However, with these systems the degree of transcription activation is limited and the regulation is not strict, as a result of which the usability for industrial applications is limited.

Furthermore, the addition of unconventional substrate components - such as cooking salt - is undesirable in production processes in the food industry, whilst temperature induction in lactic acid bacteria with, for example, temperature-sensitive repressor systems - for example as described for *E. coli* - is not always possible because lactic acid bacteria have a limited temperature range.

Other known systems for regulating the expression in lactic acid bacteria require inducers which are toxic and therefore not acceptable for foodstuffs, such as mitomycin C.

The systems described above also have the disadvantage that with these systems the regulation is generally negative regulation, that is to say repression of the gene expression during growth under specific conditions, which is less simple to use (more laborious) than the use of positive regulation by adding a specific substance at a desired point in time.

Furthermore, in cases where lactic acid bacteria are used in foodstuffs, such as, for example, in cheese and yoghurt, it is undesirable first to make substantial changes to the composition of the medium in order to make it possible to regulate the transcription of a specific gene, and this is a disadvantage of all of the systems described above.

There is therefore a need for a simple method for the controlled expression of a DNA fragment coding for one or more genes in lactic acid bacteria, with which method the above-mentioned disadvantages can be overcome, so that the transcription of the one or more desired genes can be regulated positively and to a marked degree at any desired point in the growth of the lactic acid bacteria concerned, without adversely influencing the composition of the medium and thus also the desired product characteristics of the fermented product to be obtained.

This certainly also applies in cases where lactic acid bacteria are used as production organisms for foodstuffs or constituents thereof where the production has to take place in a controlled manner.

It has now been discovered that a system of this type can be obtained by making use of promoters of microbial genes which code for antimicrobial peptides from lactic acid bacteria, such as the promoter of the nisin A structural gene.

This structural gene, as well as its promoter and the sequence thereof, have already been described in the prior art [Buchman et al.; J. Biol. chem. 263 (31) (1988) 16260-16266].

International Patent Application 93/20213 (Agriculture and Food Research Council) also describes the sequence of the *nisA* promoter and its flanking regions, as well as plasmids which contain this promoter.

However, the said application relates to organisms and methods for the production of variant nisins, including nisin Z, which in general are produced by mutating and/or completely inactivating the *nisA* structural gene.

However, the expression of these mutated nisin structural genes is preferably carried out under the influence of a *lacA* promoter; to this end the naturally occurring nisin promoter is replaced by said heterologous promoter.

However, the problem addressed by the present application, that is to say the provision of a preferably homologous expression system in *Lactococcus*, which is able to give a strict/absolute regulation of the gene expression using (in low concentrations) suitable inducers, which preferably are acceptable for food products, is not solved by the said application.

This is also apparent from a discussion of International Patent Application 93/20213 by one of its inventors, M.J. Gasson, in FEMS Microbiol. Rev. 12 (1993), 3-20, in which it is stated that

"while the lactose operon is subjected to negative regulation it is only partially switched off during repression. More effective genetic switches are desirable [...]."

J.R. van der Meer, O.P. Kuipers, W.M. de Vos et al., Journal of Bacteriology, May 1993, p. 2578-2588 describe the *nisP* and *nisR* genes of *L. lactis*. In this article it is stated that the *nisR* gene product is a "protein regulatory involved in nisin biosynthesis".

However, the *nisR* gene product is not a protein with antibacterial activity but a factor which - together with the *nisK* gene product - is involved in the signal transduction of extracellular nisin, which results in transcription activation. Moreover, although the *nisR* gene does contain a promoter, this promoter is not induced by the *nisR* gene product, nor by nisin or its derivatives. Moreover, the *nisA* and *nisZ* promoters cannot be induced by the *nisR* gene product.

It has now been discovered that promoters which code for antimicrobial peptides, and in particular the *nisA* promoter and the *nisZ* promoter, can be induced by their own gene product, that is to say the antimicrobial peptide for which they

regulate the expression in nature, and/or derivatives thereof. Thus, it has been found that the *nisA* promoter can be induced strictly/absolutely by nisin or suitable analogues/derivatives thereof, for example derivatives obtained after post-translational expression. By this means it is possible, by adding such an inducer/inducing factor in a controlled manner at a desired point in time, to obtain expression of a homologous or heterologous gene which is operably connected to said antimicrobial promoters.

The present invention therefore relates to a method of the type described above, characterised in that

- the DNA fragment is under the control of a promoter of a microbial gene which codes for an antimicrobial peptide, and
- the genes coded for on the DNA fragment are brought to expression by the addition of a suitable inducer for the transcription activation.

The promoter is generally selected from promoters of genes which code for antimicrobial peptides, such as, for example, the lantibiotics, and examples of such genes will be apparent to those skilled in the art.

Furthermore, a person skilled in the art will be able, on the basis of the description given below, to determine which promoters are suitable for use in the present invention, and which inducing factors must or can be used with these promoters.

The promoter used is preferably derived from lactic acid bacteria, more particularly from the species or strain which is used for bringing the desired DNA fragment to expression, so that the promoter is homologous with respect to said expression host.

A non-limiting example of a promoter according to the present invention, which is also preferred, is the promoter of the *nisA* gene or the *nisZ* gene from *Lactococcus lactis*.

The inducer used can be any compound which induces the gene expression. The inducing factor employed depends on the promoter used and a person skilled in the art will be able to determine, on the basis of the description given below, which inducing factor is suitable for the promoter used.

The preferred inducers - such as nisin and derivatives thereof - are advantageously acceptable for foodstuffs, more particularly for dairy products, so that the method according to the invention can be employed in the preparation of these products. The inducers used also offer a substantial advantage in the production of pharmaceuticals by lactic acid bacteria, because they are not toxic and consequently offer a relatively inexpensive option for a safe production method for these substances.

Usually, and preferably, the inducer will be an antimicrobial peptide or a suitable derivative, analogon, mutant, fragment and/or variant thereof, the inducer activity of which is sufficient to give induction. The latter inducers can be obtained, for example, by post-translational modification *in vivo*, or by preparing a derivative in a manner known per se, for example enzymatically or via a chemical route.

Furthermore, the inducer used will usually be derived from the gene product of the (structural) gene which in the wild type is under the control of the promoter used. For instance, the *nisA* or *nisZ* promoter preferably used can be induced by derivatives of nisin, which are preferably chosen from nisin A, nisin Z and/or analogues, mutants, variants and/or derivatives thereof.

The inducers are preferably produced in and by a lactic acid bacterium, so that said inducers are not alien to the lactic acid bacterium used and/or the production process used. This also makes it possible simply to add the inducer in the form of (a supernatant of) a culture of a lactic acid bacterium which produces this inducer, which offers appreciable advantages from the standpoint of process engineering.

The inducer used is therefore acceptable for foodstuffs, relatively inexpensive, simple to produce and easy to add, without having an adverse effect on the culture medium or the end product.

However, the invention is not restricted to the above-mentioned preferred inducers, but comprises, in general, the use of all factors which are able to induce the expression in a suitable manner.

The present invention is based on the fact that certain promoters of genes, especially of gene clusters, which code for antimicrobial peptides can be regulated externally in a simple manner, in particular by adding their gene product - optionally after further post-translational modifications - or a suitable derivative, analogon, mutant, fragment and/or variant thereof.

This is surprising, because the antimicrobially active inducer gives induction at levels which are far below the MIC value, the minimum inhibitory concentration (De Vos et al. (1993) Appl. Environm. Microbiol. 59, 213-218) for the micro-organisms used.

The invention therefore comprises, in the broadest terms, the culture of a lactic acid bacterium which contains a desired gene under the control of a promoter as described above. The culture of said lactic acid bacterium can be carried out in the conventional manner, in conventional culture media and using the conventional medium components.

Usually and preferably, milk, milk-containing and/or milk derived media are used. It should be noted that in general, such media contain all the substances required for the growth of the lactic acid bacteria, so that further medium components to facilitate growth of the lactic acid bacteria is not necessary, although this is included within the scope of the invention.

The induction of the expression of the desired gene can then be obtained by positive regulation, that is to say by adding the inducer to the culture described above.

The addition of said inducer can be carried out at any desired point in time during the culture and particularly at the point in time when expression of the gene is desirable.

Furthermore, advantageously only very small amounts - i.e. less than 1 µg/l - of the inducing factor will be required. Thus, in the case of the *nisA* promoter, it is sufficient to add the inducing nisin derivatives in an amount of 0.000001 - 10 mg/l, preferably 0.00001 - 0.1 mg/l, based on the total volume of the culture, for example in the form of a small amount of a fermentation fluid of a nisin-forming *L. lactis* strain.

The DNA fragment with the one or more desired genes can be brought under the control of the promoter by any means known per se, and suitable techniques will be apparent to persons skilled in the art and/or will be described in more detail below and in the abovementioned prior art, incorporated herein by reference.

In general, this comprises operably linking a nucleotide sequence coding for the one or more genes to be expressed to a nucleotide sequence coding for the promoter, the term "operably linked" having the usual meaning in the art, i.e. that the nucleotide sequences are in the correct reading frame with respect to each other, or possess their own binding sites, so as to enable translation and optionally polycistronic transcription (when the genes code for functional proteins or polypeptides) of the genes under the control of the promoter. (All other terms used in the present description, when not specifically defined herein, have the general meaning as known in the art).

In order to operably link these nucleotide sequences, the nucleotide sequence of the structural gene can be inserted into the DNA material on which the promoter is present and/or naturally occurs, optionally after inactivation, mutation or deletion of the genes naturally controlled by said promoter, the nucleotide sequence coding for the promoter can be inserted into the DNA material carrying the structural gene to be expressed, i.e. in front of the structural gene, or the nucleotide sequence coding for the promoter and the nucleotide sequence coding for the structural gene can be linked to form a new DNA material for instance an expression cassette maintained on a plasmid. With all these techniques, the nucleotide sequence coding for the promoter and/or the nucleotide sequence coding for the structural gene(s) can be identified, isolated and optionally transferred, inserted and/or linked by means of conventional genetic manipulation and cloning techniques as for instance described in the prior art, mentioned herein, which is incorporated by reference.

Furthermore, any desired DNA fragment which codes for a desired characteristic, a desired protein, enzyme or peptide, or a desired ribonucleic acid can be brought to controlled expression by the method of the invention.

When the structural gene to be expressed codes for a protein or polypeptide, this peptide can also be a precursor of the mature polypeptide, such as a pre, pro or prepro form thereof, which can undergo further post-translational modifications. As such, the invention comprises both expression of the mature protein as well as expression of these "ripening forms", with or without further post-translational modifications depending upon whether the expression host contains the cellular and enzymatical mechanisms to carry out said modifications.

The invention further comprises both secretion of the formed polypeptide or protein (in mature and/or ripening form), i.e. by crossing the bacterial cell wall and/or cell envelope. The invention also comprises intracellular accumulation of the desired

polypeptide or protein (in mature or ripening form), optionally followed by disruption of the cell wall so as to free said protein or polypeptide into the medium, i.e. by lysis or by the use of holines as described herein below.

The invention also comprises the expression of in-frame fusions of the desired protein/polypeptide, and the protein naturally controlled by the promoter, under the control of said promoter. After that, the desired protein or polypeptide can be obtained by cleavage of said fusion in a manner known per se.

In the method according to the invention, genes on said DNA fragment are preferably brought to increased expression. In this context it is generally possible to obtain an increase in the expression by a factor of more than 100, which is appreciably more than the expression ratio which can be obtained using systems from the prior art. Furthermore, in the absence of the inducing factor, a very low level of transcription for the desired gene is obtained.

The increased expression, in combination with the very strict regulation, makes the method according to the present invention particularly suitable for the overproduction of gene products at a specific point in the growth of lactic acid bacteria, and methods of this type constitute a further aspect of the invention.

The method of the invention therefore makes it possible to obtain efficient expression of desired homologous or heterologous genes in lactic acid bacteria by positive regulation of the transcription of said genes. Said regulation can be obtained by the addition of a component of a naturally occurring fermentation product, which can be added at various points in time without it being necessary first to make other modifications to the composition of the medium.

All these factors contribute to the very wide applicability of the method according to the invention, so that the present application meets a need which has already existed for a long time and thus constitutes an important step forwards in the application of regulatory gene expression systems. Further advantages of the present invention will become apparent from the description given below.

The method according to the present invention is in particular suitable for use in the preparation of proteins and/or RNA in lactic acid bacteria and methods of this type constitute a further aspect of the invention.

In this context, the method according to the invention makes it possible, in particular, to separate the growth of the production organism from the production of the desired RNA or protein. This can be very useful in the optimisation of product production on a large scale.

The method according to the present invention can be used for the preparation of proteins such as enzymes, membrane proteins, lytic proteins, extracellular proteins, antimicrobial proteins, and the like.

Thus, according to a particular aspect, the method of the invention is very suitable for the overproduction of antimicrobial peptides, such as nisin A, nisin Z and/or derivatives or mutants thereof, by bringing coding structural genes onto a plasmid under the control of the promoter fragment described. Said overproduction can be carried out using a wild type lactic acid bacterium with a conventional nisin promoter, such as that of *nisA*, and induction with nisin or a derivative thereof.

The familiarity with the inducer is of great importance for increasing the production of antimicrobial peptides, and methods for this constitute a further aspect of the invention. This can take place, for example, by achieving a sufficiently high concentration of the inducer, for example by addition, under various fermentation conditions, such as batch culture, continuous culture, or production by immobilised cells. In addition, it is possible to guarantee a sufficient degree of induction by, for example, immobilisation of the inducer.

Other desired proteins and/or gene products can be obtained by operably placing genes coding for the product of interest in the chromosome of the microorganism downstream of the promoter with the aid of well-known chromosomal integration techniques (see, for example, Leenhouts, K.J., Kok, J. and Venema, G. (1991) *J. Bact.*, 173, 4794-4798 and Kuipers, O.P., Beerthuyzen, M.M., Siezen, R.J. and de Vos, W.M. *Eur. J. Biochem.*, 216 (1993), 281-291, and Sambrook et al - also referred to as Maniatis et al - molecular cloning, A Laboratory Manual, Second edition, Cold Spring Harbor Press, 1989, the contents of which are incorporated here by reference), or by cloning the said genes in expression vectors such as plasmids on which the relevant promoter fragment is present, and then bringing said genes to expression in a lactic acid bacterium.

The method according to the present invention can, moreover, advantageously be used for the production, in a regulated manner, of RNA molecules which are active as "anti-sense RNA" with respect to a gene to be inactivated, so that specific genes can be inactivated. The possibility of the use of anti-sense RNA in lactic acid bacteria has already been described in the prior art, but was restricted by the absence of usable induction systems (see, for example, S.G. Kim and C.A. Batt, *Appl. Environm.* 57 (1991), 1109-1113.)

Furthermore, the method according to the invention can be used for the production of RNA or proteins which have a negative effect on the growth of the lactic acid bacterium used, such as lytic enzymes and other gene products which are intrinsically lethal.

According to this aspect of the invention, controlled lysis of lactic acid bacteria can be achieved, for example so as to be able to induce and to control the release of intracellular enzymes - which according to the method of the invention can be produced intracellularly - in the final fermented product. This offers great advantages for the production and accelerated maturing of various fermented foodstuffs, such as, for example, cheese and yoghurt.

An advantage in this context is, in particular, that the controlled lysis can be induced at any desired point in time during culture by the addition of the inducer, said inducer preferably being acceptable for foodstuffs.

It will be clear that such methods could not, or could hardly, be achieved with systems from the prior art because these known systems were not able to produce the strict/absolute regulation required for this purpose or were not acceptable for use in the preparation of foodstuffs.

For instance, M.J. Gasson, *FEMS Microbiol. Rev.* 12 (1993) 3-20, describes a system in which a *Listeria* bacteriophage lysine is brought to expression in *L. lactis* under the control of a *lac* promoter, in which context it is stated that:

"While the lactose operon is subject to negative regulation it is only partially switched off during repression. More effective genetic switches are desirable [...]"

With the method according to the present invention a virtually absolute positive regulation is obtained, which makes it possible to bring the lysines or other genes to expression highly selectively.

In a further aspect, the invention therefore relates to methods for the preparation of foodstuffs, in particular dairy products, in which the expression system according to the present invention is used. The preparation of these food products using the invention can be carried out analogously to known methods, for example by using a lactic acid bacterium with an expression system according to the invention, and induction of the expression by the addition of a suitable medium component which is safe in food.

It will be clear to a man skilled in the art that in the production of dairy products, the culture medium will be milk or a milk derived or milk containing medium, which is usually used for the production of the relevant dairy product.

Finally, the invention relates to a lactic acid bacterium which contains one or more homologous or heterologous genes, which

- have been brought under the control of a promoter of a microbial gene (cluster) which codes for an antimicrobial peptide, and which
- differ from the genes which are naturally - i.e. in the native form or the wild type - regulated by this promoter.

According to a specific embodiment, the promoter in this procedure is the promoter of the *nisA* or *nisZ* gene of *L. lactis* and the homologous and heterologous gene does not belong to the nisin gene cluster described below.

The one or more genes can be introduced into the chromosomal DNA of the microorganism or by means of a suitable vector, such as a plasmid, in accordance with techniques which are known per se and are described below.

In a further aspect, the invention therefore relates to a transformation vector, containing promoter of a microbial gene, which codes for an antimicrobial peptide, and one or more genes which are brought under the control of this promoter and which differ from the genes which naturally - i.e. in the native form or in the wild type of the relevant microorganism from which the promoter originates - are regulated by this promoter. The transformation vector is preferably a plasmid, but other suitable transformation vectors, such as transposons, phages and the like, can also be used and can be obtained by methods known to those skilled in the art.

The transformation vector can also be an expression vector, such as a plasmid, in which the one or more genes on the vector are expressed under the control of the promoter present on the vector.

The use of such expression vectors offers a very simple way of expressing heterologous structural genes in lactic acid bacteria, to transfer and/or incorporate a desired heterologous promoter into lactic acid bacteria, and/or both.

Therefore the promoter and the gene(s) controlled by it can be expressed from a suitable expression vector, or after incorporation of their sequences into the host's DNA.

One example of a microorganism of the invention is the lactic acid bacterium *L. lactis* NZ 9800 + pNZ 8008, which was filed with the Centraal Bureau Schimmelcultures (CBS; Central Bureau of Mould Cultures) in Baarn on 16 November 1994 under number CBS 563.94. This microorganism contains the plasmid PNZ 8008 according to the invention, comprising the *gusA* gene under the control of the *nisA* promoter of *L. lactis*.

Finally, the invention relates to the use of a lactic acid bacterium as described above in the above method(s) according to the invention.

However, the applicability of the invention is not restricted to lactic acid bacteria. Other microorganisms, in particular Gram-positive bacteria such as *B. subtilis*, can also be used in an analogous manner, as will be apparent to those skilled in the art.

The present invention is, however, of particular advantage in the preparation of foodstuffs with the aid of lactic acid bacteria.

The invention is explained in more detail below by means of a preferred embodiment, in which the promoter of the *nisA* gene from *L. lactis* is used. As already stated above, the invention is not restricted to this specific promoter.

The method according to the said preferred aspect of the present invention offers a possibility for activating the transcription of any desired gene which is placed under the control of a specific promoter fragment. In this context, use is made of a fragment containing *nisA* promoter which can be smaller than 320 bp, with a naturally occurring antimicrobial peptide as inducer.

In this context it has been found that the inducer for transcription activation of the *nisA* gene in *L. lactis* strains is the completely modified and proteolytically activated gene product of *nisA* or *nisZ*: nisin A or nisin Z (and derivatives thereof), and that this can be used as such as inducer in very low concentrations (less than 10 ng/ml).

This is surprising, partly because of the fact that to date it was known only that autoregulation can occur through the own gene product, but not that, for this purpose, post-translational modifications have to take place on this gene product before the inducing characteristic is produced. The unmodified gene product is not capable of playing the inducing role. This form of autoregulation for the production of antimicrobial peptides, in particular those which are produced by lactic acid bacteria, is a surprising and novel phenomenon.

Furthermore, neither in the case of research on subtilin - or of research on nisin - have there ever previously been indications of the need for the presence of a specific medium component in connection with regulation of gene expression of the structural gene or of other genes located in the gene cluster.

The *nisA* promoter preferably used - like the *nisZ* promoter which is also suitable - forms part of the gene cluster of *L. lactis* which is responsible for the production of nisin, a well-characterised antimicrobial peptide which is produced by various strains of the lactic acid bacterium *L. lactis* and is used in the food industry as a natural preservative because it inhibits the growth of undesirable Gram-positive bacteria (see Kuipers et al., ECB6: Proceedings of the 6th European Congress on Biotechnology, Elsevier Science, 1994).

Nisin belongs to the group of antimicrobial peptides which are referred to as 'lantibiotics'. Molecules of this type are characterised by a low molecular mass (< 4 kDa), by the presence of unsaturated amino acid residues, such as dehydroalanine and dehydrobutyrine and by the thioether bridges of lanthionine residues, which are produced by the addition of the sulfhydryl group of cysteines to the double bond of the above-mentioned unsaturated amino acids.

It is assumed that post-translational modifications of the ribosomal synthesised precursor peptide of 57 amino acids, which has no antimicrobial activity, take place, followed by secretion and the removal of the leader peptide, which leads to the active nisin in the medium.

To date, 8 genes, for which the nucleotide type sequence has been determined, which are involved in the biosynthesis, secretion, regulation and immunity of nisin have been identified. These genes are located on an approximately 12 kb fragment of the nisin/sucrose transposon Tn5276 or Tn5278 or Tn5301, and are named successively *nisABT*-



*CIPAK* or *nisZBTCIPAK*. The function of these genes in the nisin biosynthesis is only partly known. The structural gene *nisA*, and also the naturally occurring variant thereof, *nisZ*, code the unmodified precursor peptide of nisin which consists of 57 amino acid residues. The promoter sequence of the structural gene and the transcription start have been identified and a clear transcript of approximately 260 bp can be demonstrated after Northern blotting, see Kuipers, O.P., Beerthuyzen, M.M., Siezen, R.J., and de Vos, W.M., *Eur. J. Biochem.* 216 (1993), 281-291, the contents of which are incorporated here by reference. However, this article gives no information on the induction of this promoter.

It has been demonstrated that transcription of the *nisA* gene is dependent on the integrity of *nisA* itself, because a 4 bp deletion in the middle of the *nisA* gene leads to a complete absence of transcription of this truncated  $\Delta$ *nisA*, see Kuipers, O.P., Beerthuyzen, M.M., Siezen, R.J., and de Vos, W.M., *Eur. J. Biochem.* 216 (1993), 281-291, the contents of which are incorporated here by reference.

It is also known that the presence of an intact *nisR* is in all cases required for expression of *nisA*. On the basis of the existing homologies between *nisK* and *nisR* with well-characterised regulator genes belonging to the family of two-component regulator genes, it can be assumed that the *nisK* gene product (probably a histidine kinase) acts as sensor for a factor which has been unknown hitherto and that the latter activates the *nisR* gene product via a phosphorylation reaction, as a result of which transcription activation of the *nisA* gene takes place. It is therefore assumed that *nisR* and *nisK* are involved in the expression by the *nisA* promoter and play a role in the signal transduction under the influence of an extra-cellular inducing factor present.

However, the invention is not restricted to this or another explanation.

Also, although the invention is not limited thereto, applicant has found some indication that nisin producing cells in vivo stop producing nisin once they have reached the stationary phase, or at least that the amount of nisin produced is greatly reduced. Therefore, it is assumed for the preferred promoters of the nisin gene cluster of the invention that transcription/expression of the gene(s) controlled by said promoters will for the most part take place during logarithmic growth, although the invention is not limited to any particular growth phase of the expression host, and transcription/expression during for instance the stationary phase is explicitly included.

Also, although the promoters of the invention provide for a strict/absolute regulation of gene expression, a very small amount of expression of unknown etiology at an extremely low level (less than 0.1% preferably less than 0.01%), for instance due to "leakage" or to random mutations within the genetic material of the expression host, is not excluded.

Finally, although the invention is explicitly not limited thereto, it should be noted that, especially when heterologous genes or mutant genes are expressed under the control of the promoters of the invention, the expression products thus obtained (optionally after post-translational modification) will in general not, or not to any significant extent, be able to induce the promoter.

In the invention use was made of various isogenic nisin-producing and non-producing derivatives of strain *L. lactis* MG1614 (Gasson, M.J. (1993) *J. Bacteriol.* 154, 1-9).

A transconjugant was obtained which possesses the nisin/sucrose transposon Tn5276; this strain has been named *L. lactis* NZ9700. Strain *L. lactis* NZ9800 was obtained by replacement of the *nisA* gene by a truncated  $\Delta$ *nisA* gene, which contains a 4 bp deletion in position 284-287 (Kuipers et al., 1993a). This strain is no longer capable of nisin production. Expression plasmids, such as pNZ9010 and pNZ9013 which possess, respectively, the *nisA* and the *nisZ* gene under the control of the efficient *lac* promoter from the lactose operon of *L. lactis* restore the capacity for nisin A and, respectively, Z production when they are introduced into strain NZ9800 (Kuipers et al., loc cit).

Transcription analyses of the *nisA* gene in various strains, such as NZ9700 and NZ9800, were carried out as previously described (Kuipers et al., loc cit) with the aid of a radio-actively labelled *nisA* probe. Northern blotting of *L. lactis* NZ9700 RNA revealed a clear transcript of approximately 260 nucleotides (nt), whilst this was completely absent in the case of the RNA from strain NZ9800. However, when, as is described here in the invention, an amount of nisin (of between 0.01 and 1  $\mu$ g/ml) was present in the culture medium of strain NZ9800, a clear transcript of the  $\Delta$ *nisA* gene was detected. The  $\Delta$ *nisA* gene itself can never yield an active nisin molecule by the above-mentioned frame-shift mutation. Subsequent studies demonstrated that other substances were also able to produce this transcription activation. These substances include, inter alia, the completely modified precursor nisin A, and the species subtilin leader nisin Z and S5T nisin Z obtained by protein engineering techniques which have been described previously. The latter mutant contains a dehydrobutyrine in position 5. An unmodified synthetic nisin A precursor was found to be incapable of transcription activation. Other antimicrobial peptides, such as lactacin 481, subtilin, pediocin PA-1 and lactococcin A were also incapable of transcription activation. This suggests that the modified structures in the nisin molecule play a role in the inducing capacity of nisin. Disruptions of the genes *nisB* and *nisC*, which are probably involved in the modification reactions in strain NZ9700, also resulted in inability to produce nisin A, although an intact *nisA* gene is present. In this case as well it was not possible to detect any *nisA* transcript at all. However, the transcription was able to take place again when a small amount of nisin was added to the medium. In subsequent studies it was demonstrated with the aid of transcriptional fusions between the *nisA* promoter fragment and the *gusA* gene from *E. coli*, coding for  $\beta$ -glucuronidase, with its inherent ribosome binding site in NZ9800, that  $\beta$ -glucuronidase activity was detectable only when nisin (or analogues thereof, such as S3T nisin Z, S5T nisin Z or T2S nisin Z and subtilin leader nisin Z) was introduced into the medium. The mutant T2S nisin Z, which was found to possess a Dha residue in position 2, surprisingly had both a higher



activity towards various indicator organisms and better inducing characteristics than nisin A and nisin Z. The degree of *gusA* expression was found to depend on the amount of nisin used; higher expression was found with 0.028 µg/ml than with 0.014 µg/ml. It has thus been irrefutably demonstrated that it is possible, using the nisin promoter fragment, to bring a heterologous gene to expression under strict regulation.

Consequently, it is also demonstrated that it is possible to determine the nisin concentration with the aid of a reporter gene (for example the *gusA* gene), and such methods constitute a further aspect of the invention.

By cloning a fragment of  $\Phi$ US3 of *L. lactis* which has been demonstrated to be able to produce lysis, under the control of the *nisA* promoter fragment, it was also possible to achieve lysis of at least some of the cells after induction by nisin and it was possible to demonstrate that an appreciable amount of intracellular enzymes, such as, for example, aminopeptidase N (PepN) was liberated.

*L. lactis* strains of this type which show induced lysis can be very useful in acceleration of the development of taste in fermented dairy products or have industrial applications in the extraction of intracellular proteins or enzymes from *L. lactis*.

It is assumed that the autoregulation mechanism described also plays a role in the production of other antimicrobial peptides by lactic acid bacteria. In these cases, in a manner analogous to that described here for the *nis* promoter/nisin induction system, a regulated expression system can be developed with the relevant antimicrobial peptide (or analogues thereof) as inducer, and the promoter fragment of the structural gene as the element on which the regulation takes place. The expression strain used can then be the strain producing the antimicrobial peptide, in which the structural gene is switched off using methods which in themselves are well known (Leenhouts et al., and Kuipers et al., loc cit.). A plasmid with the relevant promoter fragment and gene(s) of interest can then be introduced into this lactic acid bacterium in the customary manner.

Furthermore, it is assumed that many other gene clusters exist in lactic acid bacteria which show similarities with the nisin gene cluster. In any case, homologous genes for *nisABTCPRK* have been found in the gene clusters of a few other lantibiotics, such as epidermin from *Staphylococcus epidermidis*, subtilin from *Bacillus subtilis* and pep5 from *Staphylococcus epidermidis*, which suggests a corresponding function.

Systems of this type can be used in the present invention.

In a particularly preferred aspect, the invention finally relates to a method for the extracellular production of proteins, in which method one or more genes which code for holines are brought to expression, in addition to the one or more genes which code for the protein(s) to be produced. Surprisingly, it has been found that these holines produce an increase in the secretion of these intracellular proteins, but without the cell undergoing complete lysis, as in the case of the expression of the lytic genes described above.

This has the advantage that extracellular release of the proteins produced is obtained, if appropriate after prior intracellular modifications, possibly without the cellular production mechanism for the proteins being completely lost, so that more efficient production can be achieved, in particular in continuous and/or overproduction.

This type of use of holine genes on their own - that is to say without lysine genes - has not yet been described in the prior art.

Without being restricted to any particular explanation, it is assumed that the holines increase the permeability of the cell envelope for the proteins produced by intracellular means.

Although this aspect of the method can also be used with promoters other than the promoters according to the application, it is particularly to be preferred to use the holine genes with an expression system according to the invention, so that the advantages described above are obtained.

The preferred embodiments of this system are the same as those indicated above.

Finally, the invention offers the possibility of determining the nisin concentration in a simple manner by using lactic acid bacteria in which the *nisA* promoter fragment is coupled to a promoterless reporter gene, such as the *gusA* gene of *E. coli*.

The invention will now be described with the aid of the following figures and non-limiting examples.

The figures show:

Fig. 1. Primary structure of nisin Z and the structures of the unsaturated amino acids dehydroalanine (Dha), dehydrobutyryne (Dhb), lanthionine (Ala-S-Ala) and  $\beta$ -methylanthionine (Abu-S-Ala).

Fig. 2. Northern blots of RNA from various *L. lactis* strains induced by nisin A and non-induced *L. lactis* strains, all probed with *nisA*.

Fig. 3. Northern blots of RNA from various *L. lactis* strains induced by nisin mutants or other bacteriocines and non-induced *L. lactis* strains, all probed with *nisA*.

Figs. 4a-4b. Restriction maps of the fragment from Tn5276 containing *nisA* promoter.

Figs. 5-9. Restriction maps of constructs according to the invention, i.e. pNZ8002, pNZ8003, pNZ8008, pNZ8010, pNZ8011 and pNZ8013.

Fig. 10. Coomaassie blue stained gel of extracts of strain NZ 3900 containing pNZ 8040 producing Pep N. Lanes: 1 and 5, uninduced cells; 2, induced with 0,05 µg/l nisin A; 3, induced with 0,1 µg/l nisin A; 4, induced with 0,5 µg/l nisin A; M, the molecular weight marker.

Fig. 11. The specific aminopeptidase N activity in nmol/mg.min in strain *L. lactis* NZ3900 containing pNZ8040 in the absence of nisin A and in the presence of various inducing concentrations of nisin A.

Fig. 12. Plasmid pNZ8040 containing the nisin promoter fused to the *pepN* gene

Fig. 13. Growth of *L. lactis* NZ9800 containing pNZ8011 in time, in the absence of a nisin producing culture supernatant and in the presence of nisin containing supernatant of strain NIZO R5.

## Example 1

### Transcription studies on $\Delta nisA$ of *L. lactis* NZ9800 with and without inducing agents

With the aid of Northern blotting with a radioactively labelled *nisA* probe, it has been demonstrated that no transcription of  $\Delta nisA$  takes place in strain NZ9800. Transcription does take place by complementing  $\Delta nisA$  by *nisA* or *nisZ* on pNZ9010 and pNZ9013, respectively. Here it is now first demonstrated that in strain NZ9800 transcription activation takes place by the addition of a specific amount of nisin A.

The results of Northern blots of RNA of *L. lactis* strains NZ9700 and NZ9800 cultured with and without the presence of a specific amount of nisin A are shown in Figure 2. It can clearly be seen from this figure that the transcript of the  $\Delta nisA$  gene can be demonstrated in NZ9800 only when this strain is cultured in the presence of nisin A (0.01 to 1 µg/ml) in the medium. The conditions for these experiments were identical to those described in Kuipers et al., loc cit. and incorporated herein by reference. Subsequent studies demonstrated that substances already known per se from the prior art, i.e. nisin Z, completely modified precursor nisin A, subtilin leader nisin Z, A-4D precursor nisin Z and S5T nisin Z, were able to produce induction, although in the majority of cases the peptide concentration required to obtain induction of transcription was appreciably higher than that with nisin A or Z (Fig. 3).

Cell cultures which contain other bacteriocins, such as subtilin, lacticin 481 and lactococcin A, were not capable of producing detectable transcription activation, and the same was true of a synthetically completely unmodified nisin precursor peptide (Fig. 3). It can clearly be seen from these results that only molecules which contain a modified (mutant) nisin fraction are capable of transcription activation of  $\Delta nisA$ .

## Example 2

### Controlled expression of *gusA*

To demonstrate that genes other than *nisA* or  $\Delta nisA$  can also be brought to controlled expression with the aid of the nisin promoter fragment, transcriptional fusions were made between fragments which contain the *nisA* promoter and a promoterless *gusA* gene. To this end, the large nisin promoter fragment (*Bgl*II-*Ecl*1326II: 1444 bp, Fig. 4) was first cloned on the plasmid pNZ273, which contained the *gusA* gene and a polylinker sequence just upstream of the ribosome binding site, specifically in the *Bgl*II-*Scal* (blunted) digested vector. This construct was named pNZ8003 (Fig. 5). In order to shorten the promoter fragment, the *Bgl*II-*Tth*1111 fragment (Fig. 4) was then removed, the sticky ends were blunted with the aid of the Klenow fragment of DNA polymerase I and an autoligation was carried out. The plasmid (pNZ8008, Fig. 5) thus obtained was introduced into *L. lactis* NZ9800 by means of electroporation and it was demonstrated that no  $\beta$ -glucuronidase activity was obtained on culture of the transformant in the absence of an inducing agent. However, by means of induction with extra-cellularly added nisin A or nisin Z in concentrations varying from 0.0001 to 0.1 µg/ml it was possible to obtain appreciable  $\beta$ -glucuronidase activity; see Table 1.

Table 1

<b><math>\beta</math>-glucuronidase activity of <i>L. lactis</i> NZ9800 with pNZ8008 after induction with various amounts of nisin A.</b>	
Induction amount of nisin A ( $\mu$ g/ml)	$\beta$ -glucuronidase activity in cell-free extract (arbitrary units)
0	< 0.1
0.0001	3
0.0005	18
0.0007	45
0.0070	90

It was also demonstrated that the purified mutants S3T nisin Z (contains a  $\beta$ -methyllanthionine instead of a lanthionine between residues 3 and 7) and T2S nisin Z (contains a Dha instead of a Dhb in position 2) gave clear  $\beta$ -glucuronidase activity, whilst more of the first inducer (S3T) and less of the second inducer (T2S) was required in order to obtain the same  $\beta$ -glucuronidase activity as obtained with wild-type nisin (Table 1). This demonstrates that T2S nisin Z is a better inducer than the two wild-type nisins and also demonstrates that the type of nisin and nisin concentration in the medium also have an influence on the ultimate expression level. It was then demonstrated that no induction could be obtained with preparations containing subtilin or lactococcin A. After introduction of the same plasmid (pNZ8008) into NZ9700, a nisin A producer, constitutive  $\beta$ -glucuronidase production was found, which is to be expected because constitutive nisin A is also formed here. The *gusA* gene was also cloned approximately 300 bp downstream of *nisA*, in a vector preceded by the *nis* promoter, the  $\Delta$ *nisA* fragment and part of the *nisB* gene (pNZ8002, Fig. 5). With a comparable amount of inducer, in NZ9800 this construct gave a  $\beta$ -glucuronidase activity which was substantially lower (> 20x) than that of pNZ8008, which indicates transcription attenuation, probably caused by the presence of an 'inverted repeat' sequence, located just upstream of the translation start of *nisB*. This reduced transcription level can be used to bring to expression genes for which it is considered desirable to restrict the maximum transcription level.

### Example 3

#### Controlled lysis of lactic acid bacteria

A vector with which lysis of *L. lactis* NZ9800 could be achieved in a controlled manner has been developed for the possible application of the system described above. To this end, a cassette was cloned in plasmid pNZ8010 (a derivative of pNZ8008 with a multiple cloning site for the *gusA* gene; Fig. 5), which cassette contains the genes ORF2 and *lytA* from *L. lactis* phage  $\Phi$ US3 (Platteeuw, C. and de Vos, W.M. (1992) *Gene* 118, 115-120). This construct was named pNZ8011 (Fig. 5). The ORF2 and *lytA* genes code for proteins which are, respectively, a presumed holine and a presumed amidase. The lytic effect after expression of these genes in *E. coli* or *L. lactis* is already known. The plasmid pNZ8011 was introduced into strain *L. lactis* NZ9800 and these transformants were analysed during growth in the absence or the presence of nisin A. In the absence of nisin A normal growth took place, but in the presence of nisin A the optical density of the culture decreased sharply after about 30 min, which indicates inhibition of growth and lysis of the cells. Counts of surviving cells 2, 3 or 4 hours after induction (0.014  $\mu$ g nisin A/ml) showed a reduction in viable cells by a factor of  $10^3$  compared with the situation without induction (Table 2).

Table 2

Survival of cells which are capable of bringing lytic genes to expression under the influence of nisin induction

Number of survivors (100 %) varies from  $4 \cdot 10^8$  -  $4 \cdot 10^9$  (pNZ8008),  $2 \cdot 10^8$  -  $9.7 \cdot 10^8$  (pNZ8011).

Colony-forming units <i>L. lactis</i> NZ9800		
Nisin A	pNZ8008	pNZ8011
$\mu\text{g/ml}$	%	
0	100	100
$7 \cdot 10^{-4}$	100	0.1

In a subsequent experiment it was demonstrated that when the aminopeptidase N (PepN) was overproduced at the same time in this strain by the introduction of plasmid pNZ1120 (van Alen Boerrigter, I.J., Baankreis, R. and de Vos, W.M. (1991) *Appl. Environm. Microbiol.* 265, 18499-18503), an intracellular accumulation of PepN was detected in the non-induced state without detectable PepN activity in the medium, whilst in the induced state ( $0.028 \mu\text{g}$  nisin A/ml) more than 60 % of the total aminopeptidase N (PepN) formed was found in the supernatant of the culture after 6 hours (Table 3).

The abovementioned example reveals the great advantage of this system: without making major changes to the medium, the cells can be brought to lysis, or at least to the release of intracellular enzymes into the medium, using low nisin concentrations, which is of great importance for a number of (dairy) applications.

Surprisingly, a plasmid on which only ORF2 (assumed holine) was under the control of the *nis* promoter (pNZ8013) also led to a high percentage of secreted PepN (Table 3), after transformation of said plasmid to strain NZ9800 which contained pNZ1120 (PepN). This indicates that a holine on its own is already capable of secretion of relatively large amounts of a relatively large intracellular enzyme such as PepN (95 kDa).

This could have the advantage that the cells remain metabolically active for a longer period and, consequently, could be able to continue to produce more enzyme than is the case with total lysis.

Table 3

PepN activity in the cell-free extract (cfe) and in the supernatant (sup) of NZ9800 + pNZ1120 (pepN) + pNZ8011 (ORF2-lytA) or pNZ8013 (ORF2) with and without induction by nisin A

## pNZ8011

Time (h) after induction	Induction ( $\mu\text{g/ml}$ nisa)	cfe (%)	sup (%)
2	0.03	85	15
	0	> 98	< 2.0
3	0.03	87	13
	0	> 98	< 2.0
4	0.03	85	15
	0	> 98	< 2.0
5	0.03	78	22
	0	> 98	< 2.0
6	0.03	38	62
	0	> 98	< 2.0

## pNZ8013

Time (h) after induction	Induction ( $\mu\text{g/ml}$ nisa)	cfe (%)	sup (%)
2	0.03	81	19
	0	> 98	< 2.0
3	0.03	74	26
	0	> 98	< 2.0
4	0.03	65	35
	0	> 98	< 2.0
5	0.03	48	52
	0	> 98	< 2.0
6	0.03	34	66
	0	> 98	< 2.0

**Example 4****Overexpression of *pepN* in *L. lactis* NZ3900**

- 5 The controlled expression of aminopeptidase N of *L. lactis* in *L. lactis* NZ3900 is an example of a significant overproduction of a homologous protein in *L. lactis*. This overproduction allows rapid extraction and purification of great amounts of PepN that can be used in biochemical studies. This might be useful to assess the specific contribution of this aminopeptidase N in debittering activity after the proteolytic degradation of casein in milk and for direct application of this strain in dairy fermentations.
- 10 The *pepN* gene was cloned as a 2.4-kb PCR-fragment derived from pNZ1120 (Alen-Boerrigter van, I. J., R. Baankreis and W. M. de Vos. 1991. Appl. Environ. Microbiol. 57:2555-2561.) as a template, behind the *nisA* promoter, generating pNZ8040. The primer sequences used were 5'GCAACTGCAGGAGAAGCCATGGCTGTAAAACG 3' and 5'CCTTATTCTCGAGTTGATTGTTCTATCG 3'. This plasmid was introduced in strain NZ3900 (an MG1363 derivative containing the *nisR* and *nisK* genes integrated in the *pepN* locus on the chromosome). This strain containing pNZ8040 was grown until an OD<sub>600</sub> of 0.5 was reached and
- 15 than divided in 25 ml tubes. Three tubes were induced with  $5 \cdot 10^{-4}$ ,  $1 \cdot 10^{-4}$ ,  $5 \cdot 10^{-5}$  µg/ml nisin A, respectively, and one tube was incubated without induction. After 90 minutes growth at 30°C, cells were harvested and cell free extracts were prepared. 100 µl of the extracts were mixed with equal amounts of sample buffer and 20 µl was applied to a 10% polyacrylamide-SDS gel. Fig. 10 shows the overproduction of the 95-kDa aminopeptidase N after induction (lane 2,3,4).
- 20 This result indicates that a major portion of intracellular protein is formed by the overproduced PepN protein. The cell free extracts were also used in an activity assay. PepN activity was determined at 30°C by monitoring the hydrolysis of the substrate lysyl-*p*-nitroanilide as described before (Exterkate, F. J. 1984. Appl. Environ. Microbiol. 47:177-183). Protein concentrations were determined as described previously (Bradford, M. M. 1976. Anal. Biochem. 72:248-354). Fig. 11. shows the specific activity in nmol/mg.min. of aminopeptidase N after induction with various concentrations
- 25 of nisin A. This dose-response curve shows the high production of the desired protein by induction with nisin A. A specific PepN activity of 25000 nmol/mg.min was achieved using 0.5 µg/l nisin A as an inducer, a level that has never been reached before by use of other (constitutive) promoters. This is at least a 400 fold overproduction compared to the production level of the chromosomal encoded *pepN*.

**Example 5****Controlled lysis of *L. lactis* in milk**

- This example shows the possibility to produce lytic (lethal) enzymes by expressing genes under control of the *nisA* promoter by *L. lactis* in a dairy environment. Strain *L. lactis* NZ9800 ( $\Delta$ *nisA*, lac<sup>-</sup>, prt<sup>-</sup>) containing the plasmid pNZ8011 (*lytH-lytA* cloned behind the *nisA* promoter) was grown in milk containing 0.5% casiton, 0.5% glucose and 10 µg/ml chloramphenicol. As a control strain *L. lactis* NZ9800 containing pNZ8008 (*nisA* promoter fused to the *gusA* gene) was used. While growing the strains in milk, the OD<sub>600</sub> of samples were measured after clearing 100 µl milk sample culture
- 35 with 900 µl 0.5 M Sodium-borate/10 mM EDTA. At an OD<sub>600</sub> of 0.05 the cultures were divided in different tubes and supernatant of the wild-type nisin producing strain *L. lactis* NIZO R5 ( Galesloot, T.E., and J.W. Pette. 1957. Neth. Milk Dairy J. 11:144-151) was added.
- 40 Strain NIZO R5 was also grown in milk. The pH of this overnight culture was adjusted to 6.5 by use of 1 M NaOH. Sodium citrate (2 ml) was added to clear the milk and after clearing for 15 minutes on ice the cells were centrifuged for 30 minutes. The supernatant was used for induction. 250 µl and 25 µl of supernatant was added to 25 ml of NZ9800 cultures. After
- 45 induction by NIZO R5 supernatant the growth of the strains was measured and after 2 hours the NZ9800 + pNZ8011 cells started to lyse. This lysis was caused by expression of the *lytH*, encoding the holin, and the *lytA* gene, encoding an amidase, while the NZ9800 containing pNZ8008 strain was not lysed.
- This experiment clearly demonstrates the feasibility of controlled lysis of a dairy starter strain by use of a simple fermented, food grade product i.e. a nisin producing *L. lactis* grown in milk.

**Claims**

1. Method for the controlled expression, in a lactic acid bacterium, of a DNA fragment which contains one or more genes coding for a desired characteristic, characterised in that
  - the DNA fragment is under the control of a promoter for a microbial gene which codes for an antimicrobial peptide, and
  - the one or more genes on the DNA fragment are brought to expression by the addition of a suitable inducer for the transcription activation.

2. Method according to Claim 1, characterised in that the gene or the genes on the DNA fragment are brought to increased expression.
3. Method according to Claim 1 or 2, characterised in that the promoter is the promoter of the *nisA* gene from *Lactococcus lactis*.
4. Method according to one of Claims 1-3, characterised in that the inducer is acceptable for food products.
5. Method according to one of Claims 1-4, characterised in that the inducer is produced by a lactic acid bacterium.
6. Method according to one of Claims 1-5, characterised in that the inducer is an antimicrobial peptide or an analogon, mutant, variant and/or derivative thereof.
7. Method according to one of Claims 1-6, characterised in that the inducer is chosen from nisin A, nisin Z and/or analogues, mutants, variants and/or derivatives thereof.
8. Method according to one of Claims 1-7, characterised in that the inducer is added as (a supernatant of) a culture of a lactic acid bacterium.
9. Method according to one of Claims 1-8, characterised in that the inducer is added in an amount of 0.00001-10 mg/l, preferably 0.0001-0.1 mg/l, based on the total volume of the culture.
10. Method according to one of Claims 1-9, characterised in that the DNA fragment with one or more genes is brought under the control of the promoter in a manner known per se, by operably connecting the DNA fragment to a DNA fragment containing the promoter.
11. Method according to one of Claims 1-10, characterised in that the expression of the DNA fragment results in lysis of the microorganism or the release of intracellular proteins into the medium.
12. Method according to one of Claims 1-11, characterised in that the lactic acid bacterium is a strain of *L. lactis*.
13. Method for the production of proteins or RNA in a lactic acid bacterium by expressing a DNA fragment, characterised in that the DNA fragment is expressed by means of a method according to one of Claims 1-12.
14. Method according to Claim 13, characterised in that the protein is an antimicrobial peptide which is brought to overproduction.
15. Method according to Claim 14, characterised in that the protein is nisin.
16. Method according to Claim 13, characterised in that the RNA produced is "anti-sense" RNA.
17. Method for the preparation of dairy products, which contain a desired protein, characterised in that the desired protein is obtained by application of a method according to Claim 12.
18. Dairy products obtainable by means of a method according to Claim 17.
19. Lactic acid bacterium which contains one or more homologous or heterologous genes, which:
  - have been brought under the control of a promoter of a microbial gene coding for an antimicrobial peptide, and which
  - differ from the genes which are naturally regulated by this promoter.
20. Lactic acid bacterium according to Claim 19, wherein the promoter is the *nisA* promoter of *L. lactis*.
21. Lactic acid bacterium according to Claim 19 or 20, being a strain of *L. lactis*.
22. Use of a lactic acid bacterium according to Claims 19-21 in a method according to one of Claims 1-12, 13-16 or 17.



**EP 0 712 935 A2**

23. Use of a microorganism according to one of Claims 19-21 in the determination of the concentration of an inducer in a medium.

24. Use according to Claim 23, wherein the microorganism is a microorganism according to Claims 20-21 and the inducer is nisin, an analogon, a mutant, a variant and/or a derivative thereof.

25. Vector, containing

- a promoter of a microbial gene, which codes for an antimicrobial peptide, and
- one or more genes which are brought under the control of said promoter and which differ from the genes which are naturally regulated by this promoter.

26. Vector according to Claim 25, characterised in that the promoter is the *nisA* promoter of *L. lactis*.

27. Vector according to Claim 25 or 26, being a plasmid.

28. Method for the extracellular production of gene products, such as proteins, in a microorganism by bringing a gene coding for the gene product to expression, characterised in that the permeability of the cell wall or the cell membrane of the microorganism to the gene product is increased by expressing one or more genes which code for a holine.

29. Method according to Claim 28, characterised in that the expression of the one or more genes which code for the holine is regulated by a method according to one of Claims 1-13.

fig-1




Fig - 2




Fig - 3




fig-4a




fig-4b

NAME:

FRAGMENT 1

1446 BPS DNA

S E Q U E N C E

Bgl II

```

1  AGATCTTTAA AGCAACACCT CCTAATAAAG TATGGCTGGG AGACATGACC TATACCCTA
61  CCAAAGAAGG TACCTTATAC TTAGCCGIGA ATATCGACGT TTTTTCACGT AAGATTGTAG
121  GCTGGTCAAT GCTTTCACGG ATGCAAGATA AACTGGIGAG GGATTACTTC TTACAAGCTT
181  GTGGGAAGA ACAATCCICAG CCTGGCTTGA TTGTCATAC TGATCAAGGG AGTCAATATA
241  CAAGCTCTCG TTATCAATCT ACTCTTCGTC AAGTCGGTGG TCAATCTAGC ATGAGTCGTA
301  AAGGAATCC CTATGACAAAT GCAATGATGG AGCTTTTTTA TAAGACGCTA AAGAGGGAGC
361  TTATTAATGA T6CTCATTTT GAGACAAGAG CIGAGGCTAC TCAAGAAATA TTTAATAACA
421  TTGAGACCTA TTACAAATACA AAAAGGATGC ATTCAGGCTC TGATTACAAG TCTCCAAAAG
481  ACTTTGAAAA ATATAATCT TAAATCTCT TAAATCCGIG ICTAGTTTTT CGTIGACTTT
541  CCATTATGCT TGGATTTTTT ATTGTTTAAAT TCCCTTTTTT GTATACAAGC TCGTATCTTT
601  AACAAATAT TGGCATAICG GGTTTAAAA TACTATGIGT TTTAAGAAT CTCCTAIBAG
661  ITTGACGCCA ATAACTTAGA TTAATAACAC CGTCACCTTA TTTTTAGGCA CGTTCGGCAG
721  TAACCTTATC AAAGGTATCT CGGTCAATTA GTTTCATGAT AGTATTTACT ATTTTGACTG
781  GTTTTGTGTA TTATCCAATC GTTAAATGA CAAAACAAA TAGATAAATA GATAAATATT
841  TATGGGGAGG ACAAGTGAAC TTATCATGAT TAATTGTAAA CGATTGAGTT CTGAATGTTT
901  CAAATTATGA GGAACAACAG AGTTGGACTA TTCTTTAANC GCCTCGACGA TACCATCACT
961  CTTCAATTAGC CTAAATTTAA CAAGTTAAAA TCATTAGAAT AATCTCTTTT ACAAATAATA
1021  TTTATTTAAG TTATAGTTGA CGAATATTTA ATAATTTTAT TAATATCTTG ATTTCTAGT Tth III I
1081  TCCTGAATAA TATAGAGATA GGTTTATTGA GTCCTAGACA TACTTGAATG ACCTAGTCTT
1141  ATAACATAC TGACAATAGA AACATTAAAC AATCTAAAAC AGCTTTAATT CTATCTTGAG-35
1201  AAAGTATTGG TAATAATATT ATTGTGATA ACGGAGCAT AATAACGGC TCTGATTAAA
1261  TTCTGAAGTT TGTAGATAC AATGATTTCG TTCGAAGGAA CTACANAATA AATTATAAGG RBS
1321  AGGCACTCAA AATGAGTACA AATGATTTTA ACTTGGATTI GGTAICTGTT TCGAAGAAG
1381  ATTCAGGIGC ATCACCAGC ATTACAAGTA TTTCGCIATG TACACCCCGT TGTAAACAG
1441  GAGCTC

```

nis A

Sst I

Ecl136II

fig-5




fig - 6






fig-7




fig - 1




fig-9




fig -10




fig-11




fig-12






fig-13





**THIS PAGE BLANK (USPTO)**



(19)

Eur päisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 712 935 A3**

(12)

## EUROPEAN PATENT APPLICATION

(88) Date of publication A3:  
26.11.1997 Bulletin 1997/48

(43) Date of publication A2:  
22.05.1996 Bulletin 1996/21

(21) Application number: 95203153.2

(22) Date of filing: 17.11.1995

(51) Int. Cl.<sup>6</sup>: C12N 15/74, C12N 1/21,  
A23C 19/032, A23C 23/00,  
G01N 33/04

(84) Designated Contracting States:  
BE CH DE DK ES FR GB IE IT LI NL SE

(30) Priority: 18.11.1994 NL 9401934  
18.11.1994 NL 9401935

(71) Applicant:  
STICHTING NEDERLANDS INSTITUUT  
VOOR ZUIVELONDERZOEK  
6710 BA Ede (NL)

(72) Inventors:  
- Kuipers, Oscar Paul  
NL-6713 MS Ede (NL)  
- De Vos, Willem Meindert  
NL-6721 SJ Bennekom (NL)

(74) Representative:  
de Bruijn, Leendert C. et al  
Nederlandsch Octrooibureau  
P.O. Box 29720  
2502 LS Den Haag (NL)

### (54) Method for controlling the gene expression in lactic acid bacteria

(57) The present invention relates to a method for the controlled expression, in a lactic acid bacterium, of a DNA fragment containing one or more genes coding for a desired characteristic, characterised in that

- the DNA fragment is under the control of a promoter for a microbial gene which codes for an antimicrobial peptide, and
- the gene/the genes are brought to expression on the DNA fragment by the addition of a suitable inducing factor for the transcription activation.


The promoter is preferably the *nisA* promoter from *Lactococcus lactis*.

The inducer is preferably acceptable for food products, and more preferentially is nisin or derivatives thereof.

The invention also relates to a method for the production of proteins or RNA, as well as to a method for the preparation of dairy products using the expression system according to the invention.

The invention finally relates to lactic acid bacteria and expression vectors for use in the method according to the invention.

fig -11



EP 0 712 935 A3



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 95 20 3153

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.CL.6)
Y	ENGELKE G ET AL: "Regulation of nisin biosynthesis and immunity in Lactococcus lactis 6F3." APPL ENVIRON MICROBIOL, MAR 1994, 60 (3) P814-25, UNITED STATES, XP002041574 * the whole document *	1-29	C12N15/74 C12N1/21 A23C19/032 A23C23/00 G01N33/04
Y	D.R.KIRSCH ET AL.: "The use of beta-galactosidase gene fusions to screen for antibacterial antibiotics" THE JOURNAL OF ANTIBIOTICS, vol. 2, no. 91, pages 210-217, XP002041575 * the whole document *	1-29	
Y	WO 94 00581 A (VIAGEN OY) 6 January 1994 * claim 23 *	16	
Y	WO 93 20213 A (AGRICULTURE AND FOOD RESEARCH COUNCIL) 14 October 1993 * page 24, line 30 - page 25, line 4; claims 1-15; figures 3-5 * * page 26, line 26 - page 27, line 10 *	3,7,15, 20,24,26	TECHNICAL FIELDS SEARCHED (Int.CL.6) C12N
Y	C.A.SHEARMAN ET AL.: "Autolytic lactococcus lactis expressing a lactococcal bacteriophage lysin gene" BIOTECHNOLOGY, vol. 10, pages 196-199, XP002041576 * the whole document *	11,28,29	
-/--			
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 24 September 1997	Examiner Gurdjian, D
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPO FORM 1503 (01.92) (P04C30)



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 95 20 3153

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	AKIO KURIDA ET AL.: "Molecular cloning and sequencing of the upstream region of the major Bacillus autolysin gene : a modifier exhibiting sequence homology to the major autolysin and the spold product" JOURNAL OF GENEREAL MICROBIOLOGY, vol. 138, no. 6, pages 1067-1076, XP002041577 * the whole document *	11,28,29	
A	J.ROELOF VAN DER MEER ET AL.: "Characterization of the Lactococcus lactis Nisin A opreon genes nisP, encoding a subtilisin-like serine protease involved in precursor processing, and nisR, encoding a regulatory protein involved in Nisin biosynthesis" JOURNAL OF BACTERIOLOGY, vol. 175, no. 9, pages 2578-2588, XP002041578 * the whole document *	3,5,7,9, 12-15, 17,18, 20,21, 23-27,29	
A	G.LAKSHMIDEVI ET AL.: "Molecular characterization of promoters of the Lactococcus lactis subsp. cremoris temperate bacteriophage BK-5T and identification of a phage gene implicated in the regulation of promoter activity" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 56, no. 4, pages 934-942, XP002041579 * page 934, left-hand column, paragraph 2 - right-hand column, paragraph 1 * * abstract *	1-29	
The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>24 September 1997</b>	Examiner <b>Gurdjian, D.</b>
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 150 01.12 (P0401)



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 95 20 3153

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A,D	O.P.KUIPERS ET AL.: "Characterization of the nisin gene cluster nisABTCIPR of Lactococcus lactis " EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 216, no. 1, pages 281-291, XP002041580 * the whole document *	3,7,15, 20,24,26	
A	WO 92 04451 A (GENESIT OY) 19 March 1992 * claims 1-23 *	1,19,22, 25	
A	WO 87 00203 A (THE TRUSTEES OF COLUMBIA UNIVERSITY IN THE CITY OF NEW YORK) 15 January 1987 * claims 1-20 *	1,19,22, 25,28	
P,X	KUIPERS OP ET AL: "Autoregulation of nisin biosynthesis in Lactococcus lactis by signal transduction." J BIOL CHEM, NOV 10 1995, 270 (45) P27299-304, UNITED STATES, XP002041581 * the whole document *	1-15, 17-27	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 24 September 1997	Examiner Gurdjian, D
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 150 (1.1) (P/M/01)